

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1302CXP

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 "Ask CAS" for self-help around the clock
NEWS 3 FEB 27 New STN AnaVist pricing effective March 1, 2006
NEWS 4 APR 04 STN AnaVist \$500 visualization usage credit offered
NEWS 5 MAY 10 CA/CAPLUS enhanced with 1900-1906 U.S. patent records
NEWS 6 MAY 11 KOREAPAT updates resume
NEWS 7 MAY 19 Derwent World Patents Index to be reloaded and enhanced
NEWS 8 MAY 30 IPC 8 Rolled-up Core codes added to CA/CAPLUS and
USPATFULL/USPAT2
NEWS 9 MAY 30 The F-Term thesaurus is now available in CA/CAPLUS
NEWS 10 JUN 02 The first reclassification of IPC codes now complete in
INPADOC
NEWS 11 JUN 26 TULSA/TULSA2 reloaded and enhanced with new search and
and display fields
NEWS 12 JUN 28 Price changes in full-text patent databases EPFULL and PCTFULL
NEWS 13 JUL 11 CHEMSAFE reloaded and enhanced
NEWS 14 JUL 14 FSTA enhanced with Japanese patents
NEWS 15 JUL 19 Coverage of Research Disclosure reinstated in DWPI

NEWS EXPRESS JUNE 30 CURRENT WINDOWS VERSION IS V8.01b, CURRENT
MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP),
AND CURRENT DISCOVER FILE IS DATED 26 JUNE 2006.

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS LOGIN Welcome Banner and News Items
NEWS IPC8 For general information regarding STN implementation of IPC 8
NEWS X25 X.25 communication option no longer available

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 09:08:21 ON 28 JUL 2006

=> FILE CA

COST IN U.S. DOLLARS

SINCE FILE

TOTAL

ENTRY

SESSION

FULL ESTIMATED COST

0.21

0.21

FILE 'CA' ENTERED AT 09:08:34 ON 28 JUL 2006

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 20 Jul 2006 VOL 145 ISS 5
FILE LAST UPDATED: 20 Jul 2006 (20060720/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> S viscosity and oil and chain(w)length

341977 VISCOSITY

719338 OIL

653568 CHAIN

495185 LENGTH

39250 CHAIN(W)LENGTH

L1 210 VISCOSITY AND OIL AND CHAIN(W)LENGTH

=> d l1 ti 1-210

L1 ANSWER 1 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Rheological and tribological characteristics of the synthetic lubricants derived from vegetable oils

L1 ANSWER 2 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Synthesis of biodegradable dimmer acid esters and study of their tribological characteristics

L1 ANSWER 3 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Flexible Biradicals in Liquid and Supercritical Carbon Dioxide: The Exchange Interaction, the Chain Dynamics, and a Comparison with Conventional Solvents

L1 ANSWER 4 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Nasal delivery of propranolol hydrochloride from sorbitan monostearate organogels: Preformulation study

L1 ANSWER 5 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Physical properties of oleochemical carbonates

L1 ANSWER 6 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Formation and investigation of microemulsions based on jojoba oil and nonionic surfactants

L1 ANSWER 7 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Rheology properties of glucopyranoside stabilized oil-water emulsions: effect of alkyl chain length and bulk concentration of the surfactant

L1 ANSWER 8 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Emulsification of oil in water as affected by different parameters

L1 ANSWER 9 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Chain matching effect appearing in nanoscale boundary films

L1 ANSWER 10 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI A novel method for preparing olefin oligomer synthetic lubricants

L1 ANSWER 11 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthesis of novel alkoxylated triacylglycerols and their lubricating base oil properties

L1 ANSWER 12 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Phase diagram of α -sulfonate methyl esters derived from palm stearin/alcohol/water systems

L1 ANSWER 13 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Molecular composition from diffusion or relaxation measurements

L1 ANSWER 14 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Composition for conditioning mineral fertilizers and method for conditioning

L1 ANSWER 15 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of Hydrophilically Modified Graft Polystyrene on AOT Oil-Continuous Microemulsions: Viscosifying Effects of P(S-g-PEO) as a Function of Graft Chain Length and Graft Density

L1 ANSWER 16 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Generalized relation between surface tension and viscosity: a study on pure and mixed n-alkanes

L1 ANSWER 17 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Chemical structure designing to enhance the yield stress of electrorheological fluids based on modified chitosan compounds

L1 ANSWER 18 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Correlation of the thermal stability of phospholipid-based emulsions and the microviscosity measurements using fluorescence polarization

L1 ANSWER 19 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Structure of cubic phases in ternary systems Glucopone/water/hydrocarbon

L1 ANSWER 20 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricants with non-Newtonian rheology and their degradation in line contacts

L1 ANSWER 21 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI High-performance lubricating greases containing high-viscosity poly- α -olefin base oils for rock drill bits

L1 ANSWER 22 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI EPR spin probe study of molecular ordering and dynamics in monolayers at oil/water interfaces

L1 ANSWER 23 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Influence of the fat characteristics on the physicochemical behavior of oil-in-water emulsions based on milk proteins-glycerol esters mixtures

L1 ANSWER 24 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Use of new etonogestrel esters

L1 ANSWER 25 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Alkylated naphthalenes as high-performance synthetic fluids

L1 ANSWER 26 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Properties of modified acyl glycerol emulsifiers synthesized from fatty acid mixtures

L1 ANSWER 27 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Low-trans fats for confectionery and bakery fat compositions

L1 ANSWER 28 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Water-in-oil microemulsion: Influence of co-surfactant chain- length and nature of emulsifier

L1 ANSWER 29 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Ultra-high molecular weight drag-reducing copolymers of vinyl aromatic monomers and α -olefins

L1 ANSWER 30 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Ultra high molecular weight polyolefins useful as flow improvers for hydrocarbons

L1 ANSWER 31 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthesis of a oil-soluble copolymer of atactic polypropylene grafted with long chain alkyl acrylates for use as pour point depressant additive in lubricating mineral oils

L1 ANSWER 32 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Physical properties of saturated estolides and their 2-ethylhexyl esters

L1 ANSWER 33 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Studies on some ester based polymeric lube oil additives

L1 ANSWER 34 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Upgrading triethanolamine esterquat performance to new levels

L1 ANSWER 35 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The effect of interfacial viscosity on the kinetics of formation of silver nanoparticles using water-in-oil microemulsions as nanoreactors

L1 ANSWER 36 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Solubilization of methanol and ethanol in palm oil stabilized by medium- and long-chain alkanols

L1 ANSWER 37 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI A lubricating oil composition containing an oil soluble copolymer of atactic poly-propylene grafted with long chain alkyl acrylates for use as pour point depressant

L1 ANSWER 38 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effects of alkanols on the solubilization of methanol and ethanol in red palm oil for biofuel

L1 ANSWER 39 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Environmentally friendly insecticide specific for use on tea and vegetables

L1 ANSWER 40 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The electrical conductance and viscosity of Nigerian traditional soaps in alcoholic media

L1 ANSWER 41 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical properties of a ternary system based on medium chain triglycerides/water/mixed nonionic sugar-based surfactants

L1 ANSWER 42 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Surfactant-free O/W emulsion formation of oleic acid and its esters with ultrasonic dispersion

L1 ANSWER 43 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Unity in diversity in interfacial phenomena

L1 ANSWER 44 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Final report on the safety assessment of PEG-5, -10, -16, -25, -30, and -40 soy sterol

L1 ANSWER 45 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effects of polarity of oil on the properties of emulsions prepared using nonionic surfactants

L1 ANSWER 46 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Studies on the effect of cosurfactant on the phase behavior of water in oil (CCl₄/water/tritonX-100/n-alkanol) microemulsions

L1 ANSWER 47 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Evaporation and flow properties of several hydrocarbon oils

L1 ANSWER 48 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Some Characteristics of Sugar Ester Nonionic Microemulsions in View of Possible Food Applications

L1 ANSWER 49 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Surface chemistry of new lubrication systems for high-speed spacecraft bearings

L1 ANSWER 50 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Detailed characterization of heavy alkylated benzene fluids by multiphase one- and two-dimensional NMR spectroscopy

L1 ANSWER 51 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Preparation of low viscosity synthetic oil from cracked olefins of slack wax

L1 ANSWER 52 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricants in continuous casting of aluminium. Part 1

L1 ANSWER 53 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Rheology of vegetable oil analogs and triglycerides

L1 ANSWER 54 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effects of star-shaped pour point depressants on pour point depression and viscosity reduction of crude oils and their products

L1 ANSWER 55 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Cooperativity among Molecules at Interfaces in Relation to Various Technological Processes: Effect of Chain Length on the pK_a of Fatty Acid Salt Solutions

L1 ANSWER 56 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The influence of chemical composition on the pour-point depressant properties of methacrylate copolymers used as additives for lubricating oils

L1 ANSWER 57 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Organosilicon compounds comprising polyhydroxy-functional hydrocarbon residues, their preparation and use

L1 ANSWER 58 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Characterization of a Sucrose Ester Microemulsion by Freeze Fracture Electron Micrograph and Small Angle Neutron Scattering Experiments

L1 ANSWER 59 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Surface light scattering from cationic surfactant films

L1 ANSWER 60 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polyester-polyamide tubular moldings with good pressure resistance, flexibility, and fuel-barrier property

L1 ANSWER 61 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Chain length compatibility effects in mixed surfactant systems for technological applications

L1 ANSWER 62 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Mixing of silicone oils with monolayers of nonionic surfactant at the air/water surface

L1 ANSWER 63 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Acyl lactylate index. Predicting the performance of acyl lactylates in oil-in-water systems

L1 ANSWER 64 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Prediction of biodegradability of mineral base oils from chemical composition using artificial neural networks

L1 ANSWER 65 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The importance of subangstrom distances in mixed surfactant systems for technological processes

L1 ANSWER 66 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polymeric additives for improving flow properties and viscosity index of lubricating oils

L1 ANSWER 67 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Ethylene- α -olefin copolymer rubber having high tensile strength, good heat resistance, and no bleed out of low molecular weight components

L1 ANSWER 68 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Characterization by ^{13}C n.m.r. spectroscopy of base oils produced by different processes

L1 ANSWER 69 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Manufacture of water-thinned alkyd resins for coatings with long pot life

L1 ANSWER 70 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Antioxidant sludge control additives

L1 ANSWER 71 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The influence of chemical structure on the oxidative stability of organic sulfides

L1 ANSWER 72 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Telomer oil with improved stability and lower viscosity

L1 ANSWER 73 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lipase-catalyzed synthesis of lesquerolic acid wax and diol esters and their properties

L1 ANSWER 74 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Method for manufacture of a food additive containing a fat substitute with low caloric content, the use of which lowers serum cholesterol in animals

L1 ANSWER 75 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of chain lengths of n-alcohols on the formation of single-phase microemulsions in n-heptane/n-alcohol/sodium dodecyl sulfate/water systems

L1 ANSWER 76 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Properties of W/O-type emulsions with surfactants and oils derived from oleic acid

L1 ANSWER 77 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Aggregation of unsaturated long-chain fatty alcohols in nonaqueous systems

L1 ANSWER 78 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The viscosity and ion conductivity of polydimethylsiloxane systems: 1. Chain length and ion size effects

L1 ANSWER 79 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Technical applications of galactomannans

L1 ANSWER 80 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Solubilization of methanol and triglycerides: unsaturated long-chain fatty alcohol/medium-chain alkanol mixed amphiphile systems

L1 ANSWER 81 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Microemulsification of silicone oil

L1 ANSWER 82 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The structure and activity of polyalphaolefins as pour-point depressants

L1 ANSWER 83 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Gear lubricants

L1 ANSWER 84 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Microstructure and dynamics of water in oil microemulsions

L1 ANSWER 85 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Ion mobilities and yields in x-irradiated polydimethylsiloxane oils

L1 ANSWER 86 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Low viscosity, water-stabilized extreme pressure lubricating oils

L1 ANSWER 87 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Coolant-lubricant for metalworking

L1 ANSWER 88 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Rheological properties of aqueous solutions of phosphated alkylol amides of fatty acids

L1 ANSWER 89 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The influence of the composition of the methacrylic esters mixture on the properties of polymeric additives for motor oils

L1 ANSWER 90 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Role of n-alkane components on the flow properties of the vacuum gas oil fraction from Bombay high crude oil

L1 ANSWER 91 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of nonionic surfactants on interfacial rheological properties of water-oil systems

L1 ANSWER 92 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Phase behavior and physicochemical properties of Triton X 100 and Aerosol OT induced microemulsions of hydrocarbons and water in presence of n-butanol

L1 ANSWER 93 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Improved sucrose fatty acid ester compositions and shortening and margarines made therefrom

L1 ANSWER 94 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Conformation-controlled network formation of triblock-copolymers in water-in-oil microemulsions

L1 ANSWER 95 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Methacrylate pour point depressants and compositions for lubricating oils

L1 ANSWER 96 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Flow behavior of mono- and diester lubricants

L1 ANSWER 97 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI A microbial biosurfactant: genetic engineering and applications

L1 ANSWER 98 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The stability of milk fat/water emulsions containing monoglyceride, sorbitan esters and caseinate

L1 ANSWER 99 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polybutadiene rubber compositions for tire treads

L1 ANSWER 100 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Water repellent efficacy of wax used in hardboard

L1 ANSWER 101 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Microwave-induced stabilization and destabilization of emulsions

L1 ANSWER 102 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Rheological behavior of silicon carbide whiskers in a model injection molding system

L1 ANSWER 103 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Factors affecting linear polyester formation at oil-water interface

L1 ANSWER 104 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of chain length of alkanes on water-in-oil microemulsions

L1 ANSWER 105 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Characterization of Aerosol OT-stabilized oil-in-water microemulsions using a time-resolved fluorescence method

L1 ANSWER 106 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Brominated surfactants. Brominated surfactants as emulsifiers and weighting agents: ethoxylated nonyl phenols

L1 ANSWER 107 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Waste gas purification

L1 ANSWER 108 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Formation of starch-lipid complexes by cooking extrusion

L1 ANSWER 109 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Fluorescence probe study of oil in water microemulsions

L1 ANSWER 110 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Liposome-type artificial red blood cells stabilized with carboxymethyl chitin

L1 ANSWER 111 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI A new betaine composition for cosmetics

L1 ANSWER 112 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI "Oil in water" emulsions in sodium p-octylbenzenesulfonate-pentanol-water-decane pseudoternary system

L1 ANSWER 113 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of oil viscosity on recovery processes in

relation to foam flooding

- L1 ANSWER 114 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Interfacial phenomena in foam flooding process for heavy oil recovery
- L1 ANSWER 115 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Structural description of water-in-oil microemulsions using electrical resistance
- L1 ANSWER 116 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Production and fuel characteristics of vegetable oil from oilseed crops in the Pacific Northwest
- L1 ANSWER 117 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Pyrolysis gas chromatographic characterization of poly(alkyl methacrylate) copolymer components of polymeric viscosity index improvers
- L1 ANSWER 118 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Raman spectroscopic studies on the dynamics of a molecular "probe" (methyl iodide) in hydrocarbon solvents
- L1 ANSWER 119 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Shear stability of viscosity index improvers. Mathematical interrelationship of mechanical shear degradation of polymers in solution
- L1 ANSWER 120 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Interactions between diacylglycerophosphoethanolamines and n-alkanes in monolayers and bilayers
- L1 ANSWER 121 OF 210 CA COPYRIGHT 2006 ACS on STN
TI The adjuvant activity of fatty acid esters. The role of acyl chain length and degree of saturation
- L1 ANSWER 122 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Stability of milk fat-water emulsions containing monoglycerides and sorbitan esters. 3. Effects of viscoelastic properties of oil-water interfacial films on the stability of O/W emulsions
- L1 ANSWER 123 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Effect of surfactants on wetting dynamics in solid/liquid/liquid systems
- L1 ANSWER 124 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Effect of lubricant composition on lubricating behavior in hot rolling of low carbon steel
- L1 ANSWER 125 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Effect of the chemical structure of an additive on the antiseizure and antiwear properties of an oil
- L1 ANSWER 126 OF 210 CA COPYRIGHT 2006 ACS on STN
TI The influence of the disperse phase on the stability of oil-in-water emulsions
- L1 ANSWER 127 OF 210 CA COPYRIGHT 2006 ACS on STN
TI The stability of oil-in-water emulsions containing paraffinic oils and alkyl sulfates
- L1 ANSWER 128 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Theoretical and practical questions of the structure-rheological research of ointments. Part 2. Effect of surface active components on the rheological properties
- L1 ANSWER 129 OF 210 CA COPYRIGHT 2006 ACS on STN
TI Oil recovery by viscous waterflooding

L1 ANSWER 130 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Viscosity index improvers. 2. Relations between the structure and viscosimetric properties of polymethacrylate solutions in lube oils

L1 ANSWER 131 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Activation volume of lubricating oils in viscous flow

L1 ANSWER 132 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of vehicles on percutaneous absorption. I. Characterization of oily vehicles by percutaneous absorption and trans-epidermal water loss test

L1 ANSWER 133 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Viscosity of vegetable oils

L1 ANSWER 134 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Dependence of laser induced breakdown of silicone oil on molecular weight

L1 ANSWER 135 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The loss of electrolytes from leaves treated with hydrocarbons and their derivatives

L1 ANSWER 136 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Surface activity of silicone oils in paints. 2. Effect of the concentration and solubility of the silicone oils

L1 ANSWER 137 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricant for high-speed revolving bearings, especially in precision machining

L1 ANSWER 138 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Relation between structure, properties, and stability of ointments with surface-active agents. VI. Influence of surface-active substances on drug release

L1 ANSWER 139 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Factors affecting flavor release and uptake in oil-water emulsions. II. Stirred cell studies

L1 ANSWER 140 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Oscillatory testing of o/w [oil in water] emulsions containing mixed emulsifiers of the surfactant/Long chain type. Influence of surfactant chain length

L1 ANSWER 141 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Influence of the smoke black structure upon the printing characteristics of printing ink

L1 ANSWER 142 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Application of the phase-inversion-temperature method to the emulsification of cosmetics

L1 ANSWER 143 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthetic ester lubricating oils

L1 ANSWER 144 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Starch gel modifiers from 12-hydroxystearic acid

L1 ANSWER 145 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthetic clays

L1 ANSWER 146 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of oil composition on agglomerate formation in flotation

L1 ANSWER 147 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polyether-polyester adducts for coating compositions

L1 ANSWER 148 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Modern additives for mineral oils

L1 ANSWER 149 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Dependence of the dewaxing effect on the chemical structure of the applied solvent. I. Alcohols and esters as dewaxing agents

L1 ANSWER 150 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Emulsion and power of a third solvent

L1 ANSWER 151 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of stationary phase viscosity of efficiency of gas chromatography columns

L1 ANSWER 152 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Production of carboxylic acids by the alkali fusion of 1,2-epoxycycloalkanes .

L1 ANSWER 153 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthetic ester oils for metallurgical quenching and high-temperature lubrication

L1 ANSWER 154 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Vulcanizable organopolysiloxanes

L1 ANSWER 155 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of fatty acid chain length on the rheological properties of lithium-soap pseudogels

L1 ANSWER 156 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polymer additions giving improved fuels for automobiles

L1 ANSWER 157 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Extraction of long-chain normal alkanes from petroleum fractions

L1 ANSWER 158 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Structure and properties of dispersions of sodium soaps of saturated carboxylic acids in hydrocarbon oils

L1 ANSWER 159 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating greases

L1 ANSWER 160 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Motor fuel compositions

L1 ANSWER 161 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The entanglement region and non-Newtonian polymer flow

L1 ANSWER 162 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of polymer structure on shear stability of polymer-thickened power transfer fluids

L1 ANSWER 163 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Structure of the pseudogels produced by thickening of hydrocarbon oils and with calcium and lithium soaps of saturated carboxylic acids

L1 ANSWER 164 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Relation between the fatty acid chain length and the rheological characteristics of Ca soap pseudogels

L1 ANSWER 165 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Some effects of monomer chain length on the properties of normal α -olefin polymers using an aluminum chloride catalyst

L1 ANSWER 166 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating compositions containing mixed metal salts of fatty acids and acids obtained by alkali fusion of copolymers of α -olefins and vinyl esters

L1 ANSWER 167 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Vaccine potentiation by emulsification with pure hydrocarbon compounds

L1 ANSWER 168 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Fire-resistant hydraulic fluids

L1 ANSWER 169 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Toxicological studies on hydrocarbons. IX. The aspiration hazard and toxicity of hydrocarbons and hydrocarbon mixtures

L1 ANSWER 170 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Changes of soybean carbohydrates during growth and germination. I

L1 ANSWER 171 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Investigation of the oil fractions of Anastas'ev crude

L1 ANSWER 172 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Catalyst systems for preparing polyurethan foam

L1 ANSWER 173 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Properties of homologous 2,5-di-n-alkylbenzenesulfonates

L1 ANSWER 174 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Oil- and water-repellent fluorochemical finishes for cotton

L1 ANSWER 175 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Block copolymer of a poly(aminotriazole) and an aromatic polyamide

L1 ANSWER 176 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Additives for vanadium-containing residual heating oils

L1 ANSWER 177 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Properties of the fatty acid esters of amylose

L1 ANSWER 178 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Effect of lubricant composition on friction as measured with thrust ball bearings

L1 ANSWER 179 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polymeric pour-point depressants for lubricating oils

L1 ANSWER 180 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Some high-pressure properties of model substances and petroleum hydrocarbons

L1 ANSWER 181 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI High-pressure investigations. IV. The viscosity/pressure behavior of high-molecular substances

L1 ANSWER 182 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Sodium-base lubricating greases

L1 ANSWER 183 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Physical and chemical properties of a new series of carboxylic acids, $\text{Cl}(\text{CF}_2\text{CFC1})\text{xCF}_2\text{CO}_2\text{H}$

L1 ANSWER 184 OF 210 CA COPYRIGHT 2006 ACS on STN

TI Variation of viscosity of starch paste

 L1 ANSWER 185 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Washable oil-in-water ointments. XI. Quasi-viscosity measurements in complex emulsifying ointments

 L1 ANSWER 186 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The application of high tension short circuit sparks in chemical synthesis

 L1 ANSWER 187 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Thermal decomposition of mono-2-ethylhexyl polypropylene glycol

 L1 ANSWER 188 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Pour-point depressants and viscosity-index improvers

 L1 ANSWER 189 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Correlation of detergency of polyethenoxy tallates with physical properties

 L1 ANSWER 190 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The effectiveness of several substances as plasticizers for eucolloids. I

 L1 ANSWER 191 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthetic lubricants from polyhydroxystearic acids

 L1 ANSWER 192 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Polyester pour-point depressants

 L1 ANSWER 193 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Wood starches. II. The structure of the sapwood starch of the sample

 L1 ANSWER 194 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Mineral oil containing copolymers of α,β -unsaturated dicarboxy esters with α,β -unsaturated monocarboxy acids

 L1 ANSWER 195 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating-oil additive

 L1 ANSWER 196 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Alkyl-substituted thiophenesulfonic acid salts as detergents in mineral oil compositions

 L1 ANSWER 197 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthesis of lubricating oil through alkylation of naphthalene

 L1 ANSWER 198 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Production of olefins by the cracking of Fischer-Tropsch waxes and their conversion into lubricating oils

 L1 ANSWER 199 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Aliphatic, arylaliphatic, and cyclic (C14-C20) hydrocarbons. Synthesis

 L1 ANSWER 200 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating-oil compositions containing wax-modifying agents

 L1 ANSWER 201 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Development of additives and lubricating-oil compositions

 L1 ANSWER 202 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Synthetic lubricant fluids from branched-chain diesters. Physical and chemical properties of pure diesters

 L1 ANSWER 203 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Solubilities of unvulcanized rubbers

L1 ANSWER 204 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating oils from Fischer-Tropsch olefins, using water-gas as raw material

L1 ANSWER 205 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Organosilicon polymers. II. The open chain dimethylsiloxanes with trimethylsiloxy end groups

L1 ANSWER 206 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Nature of the carbonyl groups in polyvinyl alcohol

L1 ANSWER 207 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Viscosities of solutions of Manila copals

L1 ANSWER 208 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The adhesive power of bituminous binders and the factors influencing it

L1 ANSWER 209 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI Properties of detergent solutions. X. Some further observations on electrophoretic mobilities in detergent solutions

L1 ANSWER 210 OF 210 CA COPYRIGHT 2006 ACS on STN
 TI The physical principles of lubrication, in particular for the explosion engine

=> s l1 and vegetable(w)oil
 82029 VEGETABLE
 719338 OIL
 12745 VEGETABLE(W)OIL
 L2 6 L1 AND VEGETABLE(W)OIL

=> d l2 chib,ab 1-6

L2 ANSWER 1 OF 6 CA COPYRIGHT 2006 ACS on STN
 140:144942 Influence of the fat characteristics on the physicochemical behavior of oil-in-water emulsions based on milk proteins-glycerol esters mixtures. Granger, C.; Barey, P.; Combe, N.; Veschambre, P.; Cansell, M. (Laboratoire Milieux Disperses Alimentaires: Physico-Chimie, Formulation et Vectorisation Nutritionnelle', ISTAB, Talence, F-33405, Fr.). Colloids and Surfaces, B: Biointerfaces, 32(4), 353-363 (English) 2003. CODEN: CSBBEQ. ISSN: 0927-7765. Publisher: Elsevier B.V..

AB Oil-in-water emulsions based on 10% milk protein preparation, 0.3% mono-di-glycerides (MDG) and 8% vegetable oil were prepared for models typifying ice cream formulations. Two MDG (saturated and partially unsatd.) and four fats (oleic oil, hydrogenated and refined coconut oils, refined palm oil) were chosen to investigate the interactions occurring between the oil phase, the MDG and the milk proteins. Influence of temperature (4°) and aging (24 h at 4°) was also tested. The emulsions were characterized for protein desorption, particle size distribution and rheol. properties. The dynamic surface activity of the milk proteins and the MDG at the oil-water interface was also determined. At 20°, emulsions were mostly stabilized by proteins although the protein load at the globule surface strongly depended on the emulsifier and the oil phase natures. A displacement of the proteins adsorbed at the oil droplet interface by the lipid surfactant was a consequence of the temperature decrease and/or aging step, suggesting a disruption of the interfacial protein interactions. This disruption was more or less marked depending on the physicochem. characteristics of the surfactant and the oil used (amount of crystallized matter, fatty acid chain length and unsatn. degree). In parallel, the variation of the apparent viscosity of the various emulsions upon temperature was well correlated with the solid fat content. On the whole, the results obtained suggested

that not only the surfactant mols., i.e. emulsifiers and proteins, but also the fat used in the emulsion formulation participated in the development of the interface characteristics and rheol. properties.

L2 ANSWER 2 OF 6 CA COPYRIGHT 2006 ACS on STN

138:206781 Physical properties of saturated estolides and their 2-ethylhexyl esters. Cermak, Steven C.; Isbell, Terry A. (National Center for Agricultural Utilization Research, New Crops and Processing Technology Research, Agricultural Research Service-USDA, Peoria, IL, 61604, USA). Industrial Crops and Products, 16(2), 119-127 (English) 2002. CODEN: ICRDEW. ISSN: 0926-6690. Publisher: Elsevier Science B.V..

AB Biodegradable, vegetable oil-based lubricants must have better low temperature properties before they can become widely acceptable in the marketplace. These low temperature properties are usually measured as the material's pour point, the min. temperature at which a material will still pour. Viscosity and viscosity index also provide information about a fluid's properties where a high viscosity index denotes that a fluid has little viscosity change over a wide temperature range. Oleic acid and a series of saturated fatty acids,

butyric

through stearic, were treated with 0.4 equiv of perchloric acid at either 45 or 55 °C to produce complex estolides, dimers and tetramers of fatty acids linked through the double bond and carbonyl group. Yields varied between 45 and 65% after Kugelrohr distillation The estolide number

(EN),

the average number of fatty acid units added to a base fatty acid, varied with reaction temperature as well as with the change in saturated fatty acids. The saturate-capped, oleic estolides were esterified with 2-ethylhexanol to obtain high yields of the corresponding ester. As the chain length of saturate capping material increased from C-4 to C-10, the low temperature performance of the estolide 2-ethylhexyl esters, namely pour point, decreased to -39 °C. The other mid-chain, saturated estolide 2-ethylhexyl esters C-6 through C-14 also had superior low temperature properties compared with their competitors; i.e. soy-based, synthetic-based and petroleum-based oils. The amount of oligomerization (EN) had an important role with the viscosities. Viscosity increased with higher oligomerization and the free acid estolides were generally several hundred centistokes (cSt) more viscous than the corresponding esters. The viscosity index ranged from 122 to 155 for the free acids estolides while the estolide 2-ethylhexyl esters had slightly higher indexes which ranged from 172 to 196. These new estolide esters displayed far superior low temperature properties, and were

more

suitable as a base stock for biodegradable lubricants and functional fluids than current com. materials.

L2 ANSWER 3 OF 6 CA COPYRIGHT 2006 ACS on STN

132:278396 Rheology of vegetable oil analogs and triglycerides. Geller, Daniel P.; Goodrum, John W. (Department of Biological and Agricultural Engineering, University of Georgia, Athens, GA, 30602, USA). Journal of the American Oil Chemists' Society, 77(2), 111-114 (English) 2000. CODEN: JAOCA7. ISSN: 0003-021X. Publisher: AOCS Press.

AB The rheol. properties of 2 complex mixts. of short-chain triglycerides were exptl. determined Dynamic or absolute viscosities of the mixts. were measured

for shear rates of 0.32 to 64.69 s⁻¹ at temps. between 25 and 80°C. The comps. of the mixts. were based on the oil of the plant species Cuphea viscosissima VS-320, a natural source of short-chain triglycerides. The dynamic viscosities of these mixts. were compared to those of a traditional vegetable oil (peanut oil) and diesel fuel. The results of this comparison were used to make ests. of the performance of such triglyceride mixts. as diesel fuel substitutes, since viscosity can be a key indicator of fuel performance for possible substitute diesel fuels. The crystallization temps.

of

these 2 mixts. were also determined exptl., and the effects of crystallization on fuel

performance were projected. Addnl., the dynamic viscosities of pure triglycerides from C6:0 to C18:0 at 75°C were plotted vs. chain length. These viscosities were measured at high shear rates ($>6 \text{ s}^{-1}$) where dynamic viscosity is shear-independent. An obvious trend in the relationship between triglyceride chain length and viscosity was observed. A 2nd-order regression was used to obtain an equation for this relationship. This equation was used as a model for composition dependence of viscosity. This model was applied to the viscosities of the triglyceride mixts. examined here. There was good agreement between the model and the actual, measured viscosity values determined in this study.

L2 ANSWER 4 OF 6 CA COPYRIGHT 2006 ACS on STN

126:20789 Telomer oil with improved stability and lower viscosity. Landis, Phillip S. (International Lubricants, Inc., USA). U.S. US 5567345 A 19961022, 9 pp., Cont.-in-part of U.S. 5,454,965. (English). CODEN: USXXAM. APPLICATION: US 1995-380127 19950130. PRIORITY: US 1993-108477 19930818.

AB There is disclosed a lower range viscosity telomer oil with an acid number of <20 and a viscosity range of from 5000 sus to 12,000 sus at 40°. The lower range viscosity telomer oil product further comprises no $>4\%$ polyunsatd. fatty acids and a plurality of aliphatic rings, wherein the telomer vegetable oil is made from .apprx.5% to .apprx.15% of a conjugated triglyceride oil, wherein the conjugated triglyceride oil has at least 50% of fatty acids having at least two conjugated double bonds, and from .apprx.85% to .apprx.95% of an unconjugated unsatd. vegetable oil, wherein the unconjugated unsatd. vegetable oil has from .apprx.10% to .apprx.75% of its fatty acids being polyunsatd. and having from .apprx.16 to .apprx.26 carbon atom chain length (unbranched).

L2 ANSWER 5 OF 6 CA COPYRIGHT 2006 ACS on STN

122:165388 Aggregation of unsaturated long-chain fatty alcohols in nonaqueous systems. Dunn, R. O.; Bagby, M. O. (NCAUR, USDA, Peoria, IL, 61604, USA). Journal of the American Oil Chemists' Society, 72(1), 123-30 (English) 1995. CODEN: JAOCA7. ISSN: 0003-021X. Publisher: AOCs Press.

AB Aggregation and related phenomena in nonaq. binary and ternary solns. containing unsatd. long-chain fatty alc. amphiphiles were studied. Six C18 fatty alcs. were studied: oleyl alc., elaidyl alc., linoleyl alc., elaidolinoleyl alc., linolenyl alc., and elaidolinolenyl alc. Equivalent conductivity and photon correlation spectroscopy confirmed that unsatd. long-chain fatty alcs. form large and polydisperse aggregates in MeOH. Critical micelle concentration (CMC) results showed that the degree of unsatn.

and

the configuration of the double bonds in the fatty alc. significantly influenced aggregation. Aggregation of oleyl alc. in a series of straight and branched medium-chain-length (C3-8) alkanol solvents was studied. For shorter-chained alkanols (C1-4), decreasing solvent dielec. constant decreased the CMC; however, for longer-chained alkanols (C4-8), no significant effects occurred on the CMC. The effect of solubilized soybean oil on the viscosity of long-chain fatty alc.-MeOH solns. was also analyzed. Relative viscosity results were consistent with those expected for microemulsions. Although preliminary in nature, these results generally support the notion that soybean oil is solubilized by incorporation into large soybean oil-in-fatty alc. aggregates in MeOH solvent, resembling a nonaq. detergentless microemulsion. The results have application in formulation of substitute diesel fuels.

L2 ANSWER 6 OF 6 CA COPYRIGHT 2006 ACS on STN

100:37027 Production and fuel characteristics of vegetable

oil from oilseed crops in the Pacific Northwest. Auld, D. L.; Bettis, B. L.; Peterson, C. L. (Dep. Plant, Univ. Idaho, Moscow, ID, 83843, USA). ASAE Publication (4-82, Veg. Oil Fuels), 92-100 (English) 1982. CODEN: ASPUDS. ISSN: 0197-1662.

AB The seed yield and oil production of 3 cultivars of winter rape, 2 cultivars of safflower, and 2 cultivars of sunflower were evaluated. Sunflower, oleic and linoleic safflower, and low and high erucic acid rapeseed were evaluated for fatty acid composition, energy content, viscosity, and engine performance in short-term tests. During 20-min engine tests power output, fuel economy, and thermal efficiency were compared to diesel fuel. The amount of farm-extractable oil produced from winter rape is by a factor of >2 greater than that from either safflower or sunflower. The winter rape cultivars, Norde and Jet Neuf had oil yields which averaged 1740 and 1540 L/ha, resp. Vegetable oils contained 94-95% of the kJ/L of diesel fuel, but were 11.1-17.6 times more viscous. Viscosity of the vegetable oils was closely related to fatty acid chain length and number of unsatd. bonds. During short-term engine tests all vegetable oils produced power outputs equivalent to diesel, and had thermal efficiencies 1.8-2.8% higher than diesel. Based on these results the species and cultivars of oilseed crops to be used as a source of fuel should be selected on the basis of oil yield.

=> file caold
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
95.72	95.93

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-4.26	-4.26

CA SUBSCRIBER PRICE

FILE 'CAOLD' ENTERED AT 09:10:41 ON 28 JUL 2006
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1907-1966
FILE LAST UPDATED: 01 May 1997 (19970501/UP)

This file contains CAS Registry Numbers for easy and accurate substance identification. Title keywords, authors, patent assignees, and patent information, e.g., patent numbers, are now searchable from 1907-1966. TIFF images of CA abstracts printed between 1907-1966 are available in the PAGE display formats.

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

=> S viscosity and vegetable(w)oil and chain(w)length

10180 VISCOSITY
6241 VEGETABLE
41813 OIL
249 VEGETABLE(W)OIL
6450 CHAIN
2153 LENGTH
280 CHAIN(W)LENGTH

L3 0 VISCOSITY AND VEGETABLE(W)OIL AND CHAIN(W)LENGTH

=> file fsta

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	9.74	105.67
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-4.26

FILE 'FSTA' ENTERED AT 09:11:16 ON 28 JUL 2006
 COPYRIGHT (C) 2006 International Food Information Service

FILE LAST UPDATED: 24 JUL 2006 <20060724/UP>
 FILE COVERS 1969 TO DATE.

>>> SIMULTANEOUS LEFT AND RIGHT TRUNCATION AVAILABLE IN THE BASIC
 INDEX (/BI) FIELD <<<

=> S viscosity and vegetable(w)oil and chain(w)length
 14346 VISCOSITY
 34695 VEGETABLE
 43478 OIL
 2397 VEGETABLE(W)OIL
 12045 CHAIN
 10303 LENGTH
 1252 CHAIN(W)LENGTH
 L4 0 VISCOSITY AND VEGETABLE(W)OIL AND CHAIN(W)LENGTH

=> file ca

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.86	106.53
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-4.26

FILE 'CA' ENTERED AT 09:11:29 ON 28 JUL 2006
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 20 Jul 2006 VOL 145 ISS 5
 FILE LAST UPDATED: 20 Jul 2006 (20060720/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> S viscosity and MCT
 341977 VISCOSITY
 2875 MCT
 L5 59 VISCOSITY AND MCT

=> s 15 and interesterified
683 INTERESTERIFIED
L6 1 L5 AND INTERESTERIFIED

=> d 16 cbib, ab

L6 ANSWER 1 OF 1 CA COPYRIGHT 2006 ACS on STN
141:53274 Low-residue, easy-cleaning and low-viscosity structured
lipid pan release compositions and methods. Teran, Pamela Lynn; Nakhasi,
Dilip K.; Shuman, Howard W.; Daniels, Roger L. (USA). U.S. Pat. Appl.
Publ. US 2004115332 A1 20040617, 18 pp., Cont.-in-part of U.S. Ser. No.
100,449. (English). CODEN: USXXCO. APPLICATION: US 2003-706633
20031112. PRIORITY: US 2002-2002/100449 20020318.

AB Medium chain triglyceride oils are interesterified with long
chain edible oils in order to form interesterified structured
lipids. These structured lipids find special application in food service
pan release cooking compns. and methods. The products have a relatively
low viscosity of between about 20 and about 52 cP while having a
smoke point which is especially suitable for cooking applications. They
provide
enhanced release properties, reduced darkening effects, less residue
build-up, and enhanced cleaning of griddling, cooking and baking pans,
containers and utensils.

=> s 15 and interesterify
22 INTERESTERIFY
L7 0 L5 AND INTERESTERIFY

=> d 15 ti 1-59

L5 ANSWER 1 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Comparative Evaluation of the Emulsifying Properties of
Phosphatidylcholine after Enzymatic Acyl Modification

L5 ANSWER 2 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Effect of lipid type on water-in-oil-emulsions stabilized by
phosphatidylcholine-depleted lecithin and polyglycerol polyricinoleate

L5 ANSWER 3 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Emulsion properties of batyl alcohol

L5 ANSWER 4 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Dynamic yielding, shear thinning, and stress rheology of polymer-particle
suspensions and gels

L5 ANSWER 5 OF 59 CA COPYRIGHT 2006 ACS on STN
TI A food surfactant containing lecithin and medium-chain triglycerides.

L5 ANSWER 6 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Bridging the gap between the mode coupling and the random first order
transition theories of structural relaxation in liquids

L5 ANSWER 7 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Disperse stabilization of water-soluble colorants with micellar liquid
crystals

L5 ANSWER 8 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Barrier hopping, viscous flow, and kinetic gelation in particle-polymer
suspensions

L5 ANSWER 9 OF 59 CA COPYRIGHT 2006 ACS on STN
TI Fatty acid ester compositions for wettability improvement of cocoa powder,
and cocoa powder and hot chocolate containing them

L5 ANSWER 10 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Mode coupling behavior in glass-forming liquid crystalline isopentylcyanobiphenyl

L5 ANSWER 11 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Corrosion-resistant, high-lubricity lubricants and lubricating greases, especially for steel wire ropes

L5 ANSWER 12 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Thermostable capsules containing curdlan, and process for producing the same

L5 ANSWER 13 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Water-soluble hot-melt adhesives comprising polymers containing free carboxylic acid groups and polyurethanes

L5 ANSWER 14 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Parameters with influence on the droplet size of w/o emulsions

L5 ANSWER 15 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Low-residue, easy-cleaning and low-viscosity structured lipid pan release compositions and methods

L5 ANSWER 16 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Observation of Liquid-to-Glass and Glass-to-Glass Transitions in L64/D20 Triblock Copolymer Micellar System

L5 ANSWER 17 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Viscoelasticity and rheology of depletion flocculated gels and fluids

L5 ANSWER 18 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Atomic transport in dense multicomponent metallic liquids

L5 ANSWER 19 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI From orientational glasses to structural glasses: What computer simulations have contributed to understand experiments

L5 ANSWER 20 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Atomic transport in dense, multi-component metallic liquids

L5 ANSWER 21 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Pressure and temperature dependence of viscosity and diffusion coefficients of a glassy binary mixture

L5 ANSWER 22 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Relaxation dynamics of a viscous silica melt: The intermediate scattering functions

L5 ANSWER 23 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Structural relaxation and frequency-dependent specific heat in a supercooled liquid

L5 ANSWER 24 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Improved Oil Solubilization in Oil/Water Food Grade Microemulsions in the Presence of Polyols and Ethanol

L5 ANSWER 25 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI A study of the microstructure of four-component sucrose ester microemulsions by SAXS and NMR

L5 ANSWER 26 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI The relaxation dynamics of a viscous silica melt: II The intermediate scattering functions

L5 ANSWER 27 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Viscous flow and jump dynamics in molecular supercooled liquids. II. Rotations

L5 ANSWER 28 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical characterization of a reverse micellar solution after loading with different drugs

L5 ANSWER 29 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Computer simulations of undercooled fluids and the glass transition

L5 ANSWER 30 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Some Characteristics of Sugar Ester Nonionic Microemulsions in View of Possible Food Applications

L5 ANSWER 31 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Adsorption of hydroxypropyl methylcellulose at the liquid/liquid interface and the effect on emulsion stability

L5 ANSWER 32 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Decoupling of diffusion from viscosity: difference scenario for translational and rotational motions

L5 ANSWER 33 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Viscoelasticity and generalized Stokes-Einstein relations of colloidal dispersions

L5 ANSWER 34 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Alcohol-free vegetable oil composition comprising a medium-chain triglyceride

L5 ANSWER 35 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Rheology and Dynamics of Colloidal Suspensions

L5 ANSWER 36 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Intraparticle Diffusion Limitations in the Hydrogenation of Monounsaturated Edible Oils and Their Fatty Acid Methyl Esters

L5 ANSWER 37 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Bimodality of the viscoelastic response of a dense liquid and comparison with the frictional responses at short times

L5 ANSWER 38 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI The glass transition: general scenario and crossover temperature

L5 ANSWER 39 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI The effect of oil components and homogenization conditions on the physicochemical properties and stability of parenteral fat emulsions

L5 ANSWER 40 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI The oleochemical industry in Malaysia: towards value addition-hand and body lotion from medium chain triglycerides

L5 ANSWER 41 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Toward a general description of the dynamics of glass formers

L5 ANSWER 42 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Comparative analysis of the fast dynamics in the supercooled nonfragile glass-forming liquid $\text{Na}_{0.5}\text{Li}_{0.5}\text{PO}_3$ observed by coherent neutron scattering

L5 ANSWER 43 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Decoupling of tracer diffusion from viscosity in a supercooled liquid near the glass transition

L5 ANSWER 44 OF 59 CA COPYRIGHT 2006 ACS on STN

TI Dynamics around the liquid-glass transition in poly(propylene glycol) investigated by wide-frequency-range light-scattering techniques
 L5 ANSWER 45 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Dynamics in a nonfragile glass-forming liquid
 L5 ANSWER 46 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Activated barrier crossing dynamics in slow, viscous liquids
 L5 ANSWER 47 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Application of mode-coupling theory to solvation dynamics
 L5 ANSWER 48 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Dynamics of strong and fragile glass formers and a scaling procedure for the temperature dependence of the viscosity
 L5 ANSWER 49 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Parametrization of viscosity-temperature relations of aluminosilicate melts
 L5 ANSWER 50 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI The dynamics of strong and fragile glass formers
 L5 ANSWER 51 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Dynamics of supercooled water: mode-coupling theory approach
 L5 ANSWER 52 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Signatures of the glass transition in a van der Waals liquid seen by neutrons and NMR
 L5 ANSWER 53 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Dynamic anomalies at the glass transition of organic van der Waals liquids
 L5 ANSWER 54 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Stretching, mode coupling, and the glass transition
 L5 ANSWER 55 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Functions of silicone oil in frying oil. X. Reconsideration of previous assumptions.
 L5 ANSWER 56 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Properties of agar gels containing MCT and konaame
 L5 ANSWER 57 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Perspective on the glass transition
 L5 ANSWER 58 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Properties of powder agar gels with MCT (medium chain triglyceride)
 L5 ANSWER 59 OF 59 CA COPYRIGHT 2006 ACS on STN
 TI Low temperature rheological properties of polymer treated lubricating oils

=> d 15 chib,ab 5

L5 ANSWER 5 OF 59 CA COPYRIGHT 2006 ACS on STN
 144:35584 A food surfactant containing lecithin and medium-chain triglycerides.. Nghee, Gwee Choon; Green, Terry (Pacifica Resources Sdn.Bhd., Malay.). PCT Int. Appl. WO 2005117600 A1 20051215, 12 pp.
 DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT,

TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2004-IB1810 20040603.

AB A food surfactant solution according to the invention comprises lecithin fractions in dry form dissolved in medium chain triglycerides (MCT) or C8 - C10 fatty acid esters of glycerols derived from caproic and caprylic acids. The very low viscosity of medium chain triglycerides enables higher lecithin concns. to be used and finer application spray droplets to be formed. The resultant food surfactant would even show that less of said solution would be needed during its application to achieve good coverage of the food powder particles. In addition, the low viscosity of the MCT allows the surfactant solution to migrate over the surface of the food powder particles at temps. down as low as 4°C.

=> s viscosity and coconut(w)oil

341977 VISCOSITY

20092 COCONUT

719338 OIL

12026 COCONUT(W)OIL

L8 670 VISCOSITY AND COCONUT(W)OIL

=> s viscosity and palm(w)kernal(w)oil

341977 VISCOSITY

15944 PALM

155 KERNAL

719338 OIL

14 PALM(W)KERNAL(W)OIL

L9 1 VISCOSITY AND PALM(W)KERNAL(W)OIL

=> d l9 chib,ab

L9 ANSWER 1 OF 1 CA COPYRIGHT 2006 ACS on STN

110:153084 Edible plastified dispersion. Cain, Frederick William; Day, Jennifer Irene; Jones, Malcolm Glyn; Norton, Ian Timothy (Unilever N. V., Neth.; Unilever PLC). Eur. Pat. Appl. EP 279499 A2 19880824, 13 pp. DESIGNATED STATES: R: AT, BE, CH, DE, ES, FR, GB, GR, IT, LI, NL, SE. (English). CODEN: EPXXDW. APPLICATION: EP 1988-200283 19880216. PRIORITY: GB 1987-3761 19870218; GB 1987-20542 19870901.

AB Low calorie spreads, containing proteins and/or hydrocolloid, which are stable at ambient temperature and under spreading conditions and which have favorable organoleptic qualities are prepared in an essentially conventional manner provided the aqueous phase has a low viscosity of <400 mPa-s and the amino acid residue content is <200 ppm (bases on weight of ag. phase). A spread with the described characteristics was prepared which contained 20 weight% continuous fat phase (26% hydrogenated soybean oil, 17.3% randomly interesterified palm and palm kernal oil at a ratio of 2:3, 55.2% sunflower oil, 1.5% Hymons 4404) and 30 weight% dispersed ag. phase (0.4% kappa-carrageenan, 1.8% NaCl, 97.8% distilled water). Amino acid content of ag. phase was 8 ppm; the viscosity was 3 cps.

=> d l8 ti 1-100

L8 ANSWER 1 OF 670 CA COPYRIGHT 2006 ACS on STN

TI Composition and wipe for reducing viscosity of viscoelastic bodily fluids

L8 ANSWER 2 OF 670 CA COPYRIGHT 2006 ACS on STN

TI Milking fat compositions prepared from natural products

L8 ANSWER 3 OF 670 CA COPYRIGHT 2006 ACS on STN

TI Use of sorbitol-substituted polydextrose in hair styling compositions
 L8 ANSWER 4 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Skin cleansing system comprising an anti-adherent formulation and a cationic compound
 L8 ANSWER 5 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Recovery of phenol from aqueous solution by supported liquid membrane using vegetable oils as liquid membrane
 L8 ANSWER 6 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Chocolate products and ingredients and methods for producing novel oil-in-water suspensions having reduced water activity levels
 L8 ANSWER 7 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Chocolate products and ingredients and methods for producing novel oil-in-water suspensions
 L8 ANSWER 8 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Treating agent compositions for animal fiber spinning
 L8 ANSWER 9 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Liquid transition nutrition for infants
 L8 ANSWER 10 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Storage-stable hyaluronate-containing compositions for application to mucous membrane, and method for stabilization of hyaluronates
 L8 ANSWER 11 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Lubricant for improved surface quality of cast aluminum and method
 L8 ANSWER 12 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Method for producing a food that contains a vegetable fat carrier, a stabilizer or a gelatinizing agent and appropriate solvents.
 L8 ANSWER 13 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Effect of coconut oil-blended fuels on diesel engine wear and lubrication
 L8 ANSWER 14 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Manufacture of biodiesel as a type of alternative fuel: transesterification of coconut oil by methanol with NaOH as a catalyst
 L8 ANSWER 15 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Hair treatment compositions containing combinations of polymers
 L8 ANSWER 16 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Hair care compositions
 L8 ANSWER 17 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Production of vegetable oil-based polyester-polyols suitable for production of stiff polyurethane foams
 L8 ANSWER 18 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Rheology of bio-edible oils according to several rheological models and its potential as hydraulic fluid
 L8 ANSWER 19 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating oil composition for internal combustion engines
 L8 ANSWER 20 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Cleaning compositions showing low temperature-dependency of viscosity

L8 ANSWER 21 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Flowable food topping compositions and methods of making and using same

L8 ANSWER 22 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Viscous fat cyclodextrin-containing compositions having low amounts of trans-fat, methods and products

L8 ANSWER 23 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Particles for the delivery of active agents

L8 ANSWER 24 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Polymer-coated particles for the delivery of active agents

L8 ANSWER 25 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Liquid cleanser compositions

L8 ANSWER 26 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Dispensing system for spraying non-fluid or viscous hair preparations

L8 ANSWER 27 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Surfactant compositions with good low temperature stability, flexibility, foamability, and moisture retaining property

L8 ANSWER 28 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Manufacture of well control fluid for workover of negative pressure well

L8 ANSWER 29 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Method and apparatus for producing edible fat-based shell for confectioneries and confectioneries produced thereby

L8 ANSWER 30 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Edible fat-based shell for confectioneries and method for producing same

L8 ANSWER 31 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Thickening agent compositions with good use of feeling useful for cleaning agents and cosmetics

L8 ANSWER 32 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Polyurethane dull white finishing paints for woodenware

L8 ANSWER 33 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Environmentally-friendly type clear paraffin control thinning agent

L8 ANSWER 34 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Vegetable oil-containing preparations for application to mucous membranes, and viscosity stabilization and feel improvement

L8 ANSWER 35 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Study of three systems of ozonized coconut oil

L8 ANSWER 36 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical behavior of oil-in-water emulsions: influence of milk protein mixtures, glycerol ester mixtures and fat characteristics

L8 ANSWER 37 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Additive composition for reclaimed pavement material

L8 ANSWER 38 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Additive composition for reclaimed pavement material

L8 ANSWER 39 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Cream substitutes comprising milk and vegetable fat and biopolymer thickener

L8 ANSWER 40 OF 670 CA COPYRIGHT 2006 ACS on STN

TI Viscous shower gel containing polyamide gelling agents and oils

L8 ANSWER 41 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent products comprising moisturizing and lubricating compositions

L8 ANSWER 42 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Properties of emulsion for cold rolling aluminum

L8 ANSWER 43 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Leave-on hair care oil-in-water emulsion

L8 ANSWER 44 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent product with improved liner treatment

L8 ANSWER 45 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Mascara composition with a keratin conditioning agent

L8 ANSWER 46 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Study on the characteristics of palm oil and its derivatives as liquid insulating materials

L8 ANSWER 47 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Rheological analysis of the structural properties effecting the percutaneous absorption and stability in pharmaceutical organogels

L8 ANSWER 48 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Hair preparations containing fluorescent nanoparticle compositions

L8 ANSWER 49 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Nature-based emulsifiers and their cosmetic applications

L8 ANSWER 50 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Milk based products containing coated alginates

L8 ANSWER 51 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Sprayable cookware release composition with fractionated oil and method of preparing food item

L8 ANSWER 52 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Cleaning compositions with good detergency, foamability, and conditioning effect

L8 ANSWER 53 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Agent-encapsulating micro- and nanoparticles, methods for preparation of same and products containing same

L8 ANSWER 54 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Low molecular weight carboxyalkyl cellulose esters and their use as low viscosity binders and modifiers in coating compositions

L8 ANSWER 55 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Crankcase lubricating oils containing ashless friction modifiers for improving fuel economy

L8 ANSWER 56 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Low molecular weight cellulose mixed esters and their use as low viscosity binders and modifiers in coating compositions

L8 ANSWER 57 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Blending of edible plant oils to obtain nutritionally favourable fatty acid ratios

L8 ANSWER 58 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Oily wax matrix suspension formulation comprising pharmacologically active agents

L8 ANSWER 59 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Hydrophilic compositions for use on absorbent articles to enhance skin barrier

L8 ANSWER 60 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Oily composition based on lipoperoxides usable in the treatment of xerostomia

L8 ANSWER 61 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Oil in water triglyceride emulsions for cream substitutes.

L8 ANSWER 62 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Low-residue, easy-cleaning and low-viscosity structured lipid pan release compositions and methods

L8 ANSWER 63 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Compositions comprising a dispersant and an overbased carboxylate or sulfonate for rheology control in coatings and inks

L8 ANSWER 64 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Storage-stable ceramide-containing emulsions and their manufacture

L8 ANSWER 65 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Hair-conditioning compositions containing polyol condensates, oils, amino-containing silicones, ethoxylated nonionic surfactants, and aliphatic alcohols

L8 ANSWER 66 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Two-phase oily cosmetic compositions

L8 ANSWER 67 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Foaming oil-in-water emulsion fat and oil compositions

L8 ANSWER 68 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Preparation and use of concentrated and ready-to-use creamer compns.

L8 ANSWER 69 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Ximenynic acid effect on food properties and health

L8 ANSWER 70 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Ximenynic acid effect on food properties and health

L8 ANSWER 71 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Moisturizing and lubricating composition

L8 ANSWER 72 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Crab core distilled oil-containing cleaning composition and its production process

L8 ANSWER 73 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Influence of the fat characteristics on the physicochemical behavior of oil-in-water emulsions based on milk proteins-glycerol esters mixtures

L8 ANSWER 74 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Mask composition containing emulsified liquid composition

L8 ANSWER 75 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Novel wax for hot melt adhesive applications

L8 ANSWER 76 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Absorbent articles with compositions for reducing irritation response

L8 ANSWER 77 OF 670 CA COPYRIGHT 2006 ACS on STN
 TI Process of making a frozen whipped topping

L8 ANSWER 78 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Fluid oil compositions with high phosphatidylserine content, their manufacture, and food compositions containing them

L8 ANSWER 79 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Oil compositions with low viscosity and coating of foods using them

L8 ANSWER 80 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Effects of oils and pharmaceutical excipients on the bioavailability of ampicillin orally administered, different oily and aqueous suspensions in rabbit

L8 ANSWER 81 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Anti-regurgitation infant formula and uses

L8 ANSWER 82 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Spinnerets apparatus for spinning spider silk fibers produced in mammalian cells

L8 ANSWER 83 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent articles with silicone elastomer-containing compositions having even distribution

L8 ANSWER 84 OF 670 CA COPYRIGHT 2006 ACS on STN
TI System for improving skin health of absorbent article wearers

L8 ANSWER 85 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Cleaning products based on microemulsions that contain oil

L8 ANSWER 86 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Comparative studies on physical properties of vegetable oils and their blends after frying

L8 ANSWER 87 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Fruit-vegetable juice and soy protein beverage and uses thereof

L8 ANSWER 88 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Edible water-in-oil emulsion with calcium

L8 ANSWER 89 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent tissues providing skin barrier enhancement

L8 ANSWER 90 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Compositions containing salts of analgesics in oil

L8 ANSWER 91 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Preparation of sugar-free chewy products and protein-based chewy products

L8 ANSWER 92 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Continuous or bicontinuous fat emulsions

L8 ANSWER 93 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Lamellar liquid crystals in viscous oil product

L8 ANSWER 94 OF 670 CA COPYRIGHT 2006 ACS on STN
TI The effects of some physico-chemical factors and pharmaceutical excipients on the bioavailability of nitrofurantoin oily and aqueous suspensions in rats

L8 ANSWER 95 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Silicone emulsion compositions with good storage stability and cosmetic materials therewith

L8 ANSWER 96 OF 670 CA COPYRIGHT 2006 ACS on STN
TI A biodegradable release agent increasing the life of concrete forms

L8 ANSWER 97 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Pharmaceutical compositions containing NO-releasing NSAID and surfactants

L8 ANSWER 98 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Non-irritating formulation for transdermal drug delivery

L8 ANSWER 99 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent articles with hydrophilic compositions containing plant extracts

L8 ANSWER 100 OF 670 CA COPYRIGHT 2006 ACS on STN
TI Absorbent articles with hydrophilic compositions containing anionic polymers

=> s 18 not emulsions
122111 EMULSIONS
L10 593 L8 NOT EMULSIONS

=> s 110 not lubricants
71972 LUBRICANTS
L11 528 L10 NOT LUBRICANTS

=> s 18 and property
571775 PROPERTY
L12 29 L8 AND PROPERTY

=> s 18 and properties
4190389 PROPERTIES
L13 134 L8 AND PROPERTIES

=> d 112 ti 1-5

L12 ANSWER 1 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Rheology of bio-edible oils according to several rheological models and its potential as hydraulic fluid

L12 ANSWER 2 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Surfactant compositions with good low temperature stability, flexibility, foamability, and moisture retaining property

L12 ANSWER 3 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Properties of emulsion for cold rolling aluminum

L12 ANSWER 4 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Hair-conditioning compositions containing polyol condensates, oils, amino-containing silicones, ethoxylated nonionic surfactants, and aliphatic alcohols

L12 ANSWER 5 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Water-borne coating composition and forming smooth multilayer coating film

=> s 112 not emulsions
122111 EMULSIONS
L14 27 L12 NOT EMULSIONS

=> d 112 ti 1-29

L12 ANSWER 1 OF 29 CA COPYRIGHT 2006 ACS on STN
TI Rheology of bio-edible oils according to several rheological models and its potential as hydraulic fluid

L12 ANSWER 2 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Surfactant compositions with good low temperature stability, flexibility, foamability, and moisture retaining property

L12 ANSWER 3 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Properties of emulsion for cold rolling aluminum

L12 ANSWER 4 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Hair-conditioning compositions containing polyol condensates, oils, amino-containing silicones, ethoxylated nonionic surfactants, and aliphatic alcohols

L12 ANSWER 5 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Water-borne coating composition and forming smooth multilayer coating film

L12 ANSWER 6 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Odorless water-in-oil emulsion inks for stencil printing

L12 ANSWER 7 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Odorless water-in-oil emulsion inks for stencil printing

L12 ANSWER 8 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI The physicochemical properties of some vegetable oils

L12 ANSWER 9 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical characterization of oils from some Kenyan plants

L12 ANSWER 10 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Effect of liquid-phase properties on ultrasound intensity and cavitation activity

L12 ANSWER 11 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Stable coconut cream alternative

L12 ANSWER 12 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Ultrasonic studies of palm oil and other vegetable oils

L12 ANSWER 13 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Effect of docosahexaenoic acid on mouse mitochondrial membrane properties

L12 ANSWER 14 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Liquid softener composition with softness, antistatic property, and water absorption for clothes

L12 ANSWER 15 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Mechanical properties of vegetable oils and fats

L12 ANSWER 16 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Relationship between the molecular structure and deep fat frying properties of edible oils

L12 ANSWER 17 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Physical and textural characteristics of some North American shortenings

L12 ANSWER 18 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Pigment dispersants with good compatibility with coating vehicle resins and manufacture thereof

L12 ANSWER 19 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Anionic liquid detergent compositions with good foaming in hard water

L12 ANSWER 20 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Interrelationships among the properties of fatty oils

L12 ANSWER 21 OF 29 CA COPYRIGHT 2006 ACS on STN

TI Rheological properties of emulsified fats with regard to their optimum consistency

L12 ANSWER 22 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Physical properties of oils and mixtures of oils

L12 ANSWER 23 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI High solids alkyd resins

L12 ANSWER 24 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Properties of dispersion polyacrylates

L12 ANSWER 25 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI s-Triazine coating resins. V. Preparation of butylated guanamine coating resins and their properties

L12 ANSWER 26 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical properties of monoglyceride sulfonate-water systems near the critical micelle concentration

L12 ANSWER 27 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Polyester pour-point depressants

L12 ANSWER 28 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI The static friction of lubricated surfaces

L12 ANSWER 29 OF 29 CA COPYRIGHT 2006 ACS on STN
 TI Some properties of sodium naphthenate used in the soap industry

=> d l13 ti 1-134

L13 ANSWER 1 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Composition and wipe for reducing viscosity of viscoelastic bodily fluids

L13 ANSWER 2 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Manufacture of biodiesel as a type of alternative fuel: transesterification of coconut oil by methanol with NaOH as a catalyst

L13 ANSWER 3 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Rheology of bio-edible oils according to several rheological models and its potential as hydraulic fluid

L13 ANSWER 4 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Cleaning compositions showing low temperature-dependency of viscosity

L13 ANSWER 5 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Viscous fat cyclodextrin-containing compositions having low amounts of trans-fat, methods and products

L13 ANSWER 6 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical behavior of oil-in-water emulsions: influence of milk protein mixtures, glycerol ester mixtures and fat characteristics

L13 ANSWER 7 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Properties of emulsion for cold rolling aluminum

L13 ANSWER 8 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Study on the characteristics of palm oil and its derivatives as liquid insulating materials

L13 ANSWER 9 OF 134 CA COPYRIGHT 2006 ACS on STN

TI Rheological analysis of the structural properties effecting the percutaneous absorption and stability in pharmaceutical organogels

L13 ANSWER 10 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Low-residue, easy-cleaning and low-viscosity structured lipid pan release compositions and methods

L13 ANSWER 11 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Foaming oil-in-water emulsion fat and oil compositions

L13 ANSWER 12 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Ximenynic acid effect on food properties and health

L13 ANSWER 13 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Ximenynic acid effect on food properties and health

L13 ANSWER 14 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Influence of the fat characteristics on the physicochemical behavior of oil-in-water emulsions based on milk proteins-glycerol esters mixtures

L13 ANSWER 15 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Spinnerets apparatus for spinning spider silk fibers produced in mammalian cells

L13 ANSWER 16 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Absorbent articles with silicone elastomer-containing compositions having even distribution

L13 ANSWER 17 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Comparative studies on physical properties of vegetable oils and their blends after frying

L13 ANSWER 18 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Lamellar liquid crystals in viscous oil product

L13 ANSWER 19 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Absorbent articles with simplified emollient compositions having good stability

L13 ANSWER 20 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Frozen slush liquid concentrate and method of making same

L13 ANSWER 21 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Controlled release encapsulated substances

L13 ANSWER 22 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Oil components modulate physical characteristics and function of the natural oil emulsions as drug or gene delivery system

L13 ANSWER 23 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Thickened oil compositions of edible oil

L13 ANSWER 24 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Lipase-catalysed production of biodiesel fuel from some Nigerian lauric oils

L13 ANSWER 25 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Preparation of redispersible dry emulsions by spray drying

L13 ANSWER 26 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI An experimental study to evaluate the use of coconut-based fuels as alternatives to diesel oil

L13 ANSWER 27 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Effect of starch-lipids inclusion complex formation on functional

properties of flour in tandoori roti

- L13 ANSWER 28 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Density and viscosity of vegetable oils
- L13 ANSWER 29 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Odorless water-in-oil emulsion inks for stencil printing
- L13 ANSWER 30 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Odorless water-in-oil emulsion inks for stencil printing
- L13 ANSWER 31 OF 134 CA COPYRIGHT 2006 ACS on STN
TI The physicochemical properties of some vegetable oils
- L13 ANSWER 32 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Enzymic interesterification of blends of castor oil and some oils rich in saturated fatty acids
- L13 ANSWER 33 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Physicochemical characterization of oils from some Kenyan plants
- L13 ANSWER 34 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Ultrasonic studies on edible oils
- L13 ANSWER 35 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Effect of liquid-phase properties on ultrasound intensity and cavitation activity
- L13 ANSWER 36 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Aliphatic acid-modified polyester polyol compositions with low viscosity and high solubility to hydrofluorocarbons
- L13 ANSWER 37 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Liquid personal cleansing compositions containing low viscosity oils pre-thickened by non-antifoaming hydrophobic polymers
- L13 ANSWER 38 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Personal cleansing bar compositions comprising low viscosity oils pre-thickened by non-antifoaming hydrophobic polymers
- L13 ANSWER 39 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Stable coconut cream alternative
- L13 ANSWER 40 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Downhole well lubricant for release of stuck coiled tubing by formation of greasy lubricating emulsion upon rubbing
- L13 ANSWER 41 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Ultrasonic studies of palm oil and other vegetable oils
- L13 ANSWER 42 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Properties of methyl esters of interesterified triacylglycerols
- L13 ANSWER 43 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Effect of docosahexaenoic acid on mouse mitochondrial membrane properties
- L13 ANSWER 44 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Rubber compositions and tire treads therefrom
- L13 ANSWER 45 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Oil-and-fat feedstock for producing frozen confections and frozen confections using the same
- L13 ANSWER 46 OF 134 CA COPYRIGHT 2006 ACS on STN

TI Low density ready-to-spread frosting and method of preparation

L13 ANSWER 47 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Mechanical properties of vegetable oils and fats

L13 ANSWER 48 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Ball point pen water-thinned ink compositions with good continuous writing properties

L13 ANSWER 49 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Oil dispersion of alumina for tape casting

L13 ANSWER 50 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Manufacture of alkyd resins with improved physico-chemical and film-forming properties

L13 ANSWER 51 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Relationship between the molecular structure and deep fat frying properties of edible oils

L13 ANSWER 52 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Preparation and use of some emulsifiers in ice cream manufacture

L13 ANSWER 53 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Viscosities of vegetable oils and fatty acids

L13 ANSWER 54 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Effect of vegetable oils on plain ice cream properties

L13 ANSWER 55 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Statistical relationships between some properties of ice cream as affected by substitution of milk fat by vegetable oils

L13 ANSWER 56 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physical and textural characteristics of some North American shortenings

L13 ANSWER 57 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Studies on viscoelastic behavior of fluid coatings. (I). Viscoelastic properties and applicability involving inter-roll transference of paints for coil coating

L13 ANSWER 58 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI New thickening agents for surfactants

L13 ANSWER 59 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Lubricating oil composition

L13 ANSWER 60 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Interrelationships among the properties of fatty oils

L13 ANSWER 61 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Edible plastified dispersion

L13 ANSWER 62 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Methyl esters of fatty acids as pesticide formulation and application aids

L13 ANSWER 63 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Rheological properties of neat household soap

L13 ANSWER 64 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Curable urethane composition

L13 ANSWER 65 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Polyester-modified vinyl resin and its use

L13 ANSWER 66 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Rheological properties of emulsified fats with regard to their optimum consistency

L13 ANSWER 67 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physical properties of oils and mixtures of oils

L13 ANSWER 68 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Effect of different suppository bases on release of indomethacin

L13 ANSWER 69 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Effect of lipid constituents on the amylograph characteristics of barley flour

L13 ANSWER 70 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI High solids alkyd resins

L13 ANSWER 71 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Alkyd coating compositions

L13 ANSWER 72 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Paper printing ink and its use

L13 ANSWER 73 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Mathematical study of a ternary system: sodium dodecyl-dioxyethyl ether sulphate, coconut oil acid amidopropyl betaine and coconut oil acid diethanol amide

L13 ANSWER 74 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Resin composition for release papers

L13 ANSWER 75 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI A spin finish for synthetic fibers and methods of lubricating synthetic yarns with an aqueous emulsion containing this spin finish

L13 ANSWER 76 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Emulsifying properties of whey protein

L13 ANSWER 77 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Water-soluble triglyceride compositions

L13 ANSWER 78 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Binder for road-marking pigments

L13 ANSWER 79 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Combination bubble-bath preparations. II. Rheological and functional effects of adjuvant surfactants in combination with sodium laurylethoxy sulfate solutions

L13 ANSWER 80 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Crystallization inhibitor for paraffin

L13 ANSWER 81 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Polyurethane foam moldings

L13 ANSWER 82 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Surface modification of fillers and reinforcement in plastics

L13 ANSWER 83 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Water soluble triglyceride compositions and method for their preparation

L13 ANSWER 84 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Effect of type of fat on starch pastes containing glycerol monostearate

L13 ANSWER 85 OF 134 CA COPYRIGHT 2006 ACS on STN

TI Composition for treating leather

L13 ANSWER 86 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Powdery, water-soluble, linear polymers or mixed polymers with very high molecular weights

L13 ANSWER 87 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical and dermatological evaluations of a Polish shampoo

L13 ANSWER 88 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Properties of dispersion polyacrylates

L13 ANSWER 89 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Factors affecting flavor release and uptake in oil-water emulsions. II. Stirred cell studies

L13 ANSWER 90 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Nutrition and biochemistry of the coconut

L13 ANSWER 91 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Nonaqueous thixotropic emulsions of emulsifiable oils and nonoily materials for use as fuels and propellants

L13 ANSWER 92 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Aerosol foam composition suitable for dispensing when warm

L13 ANSWER 93 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI s-Triazine coating resins. V. Preparation of butylated guanamine coating resins and their properties

L13 ANSWER 94 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Soldering paste

L13 ANSWER 95 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Experiments with fatty alcohol ether sulfates

L13 ANSWER 96 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Water-soluble polyester coatings

L13 ANSWER 97 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Liquid shampoo

L13 ANSWER 98 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Stability of titanium dioxide and copper phthaloxyanine pigments dispersed in alkyd-melamine resin solutions

L13 ANSWER 99 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Hard butter. I. Aspects of the rheology of fats for chocolate

L13 ANSWER 100 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Continuous viscosity measurement in stirred vessels

L13 ANSWER 101 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Photochromic compositions and substrate coated therewith

L13 ANSWER 102 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Alkyd resins

L13 ANSWER 103 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physicochemical properties of monoglyceride sulfonate-water systems near the critical micelle concentration

L13 ANSWER 104 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Aerosol topping

L13 ANSWER 105 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Effects of sucroglycerides on flour constituents and on fatty materials

L13 ANSWER 106 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Resinous coatings containing N-(1,1-dimethyl-2-hydroxymethyl)-N,N-dimethylammonium p-toluenesulfonate

L13 ANSWER 107 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Fat preparation

L13 ANSWER 108 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Hydroxy-ester modified polysiloxane resins reacted with organopolyisocyanates

L13 ANSWER 109 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Lubricating greases containing finely divided alkaline earth sulfates

L13 ANSWER 110 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Hydrogenated polyisobutylene lubricant

L13 ANSWER 111 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Edible emulsions

L13 ANSWER 112 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Preparation of liquid food for infants

L13 ANSWER 113 OF 134 CA COPYRIGHT 2006 ACS on STN
TI The stability of oils and fats after prolonged storage in retail packs

L13 ANSWER 114 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Sulfonation products

L13 ANSWER 115 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Grease hardening agents

L13 ANSWER 116 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Coating compositions containing modified urea resins

L13 ANSWER 117 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Methyl methacrylate polymers and coating compositions therefrom

L13 ANSWER 118 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Cosmetic detergent compositions

L13 ANSWER 119 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Dry-cleaning compositions

L13 ANSWER 120 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Lubricating greases containing soap-salt thickeners

L13 ANSWER 121 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Effects of foaming catalysts on hydrolysis-aging of urethan foams

L13 ANSWER 122 OF 134 CA COPYRIGHT 2006 ACS on STN
TI β -Aminopropionate surfactants

L13 ANSWER 123 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Semimicro butyric acid values, semimicro total values, and semimicro residual values of cocoa butter and cocoa-butter-substitute fats

L13 ANSWER 124 OF 134 CA COPYRIGHT 2006 ACS on STN
TI Lubricating-oil additive

L13 ANSWER 125 OF 134 CA COPYRIGHT 2006 ACS on STN
TI The relation of fats to the texture, crumb and volume of bread

L13 ANSWER 126 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Fat-soluble condensation products

L13 ANSWER 127 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Chemical constants and reaction products during development of tallowiness
 [in fats]

L13 ANSWER 128 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI The physical properties of mixtures of coconut and castor oil

L13 ANSWER 129 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI The static friction of lubricated surfaces

L13 ANSWER 130 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Physical properties of some Philippine vegetable oils

L13 ANSWER 131 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Mosquito oil and larvicide specifications

L13 ANSWER 132 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI The structural properties of anisotropic solutions of soap as
 determined by a new centrifugal falling-ball method

L13 ANSWER 133 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Some properties of sodium naphthenate used in the soap industry

L13 ANSWER 134 OF 134 CA COPYRIGHT 2006 ACS on STN
 TI Blended fatty food products

=> file stnguide

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	148.80	255.33
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-2.13	-6.39

FILE 'STNGUIDE' ENTERED AT 09:20:35 ON 28 JUL 2006
 USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
 COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
 AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
 LAST RELOADED: Jul 21, 2006 (20060721/UP).

=> file ca

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.66	255.99
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-6.39

FILE 'CA' ENTERED AT 09:27:03 ON 28 JUL 2006
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2006 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is
 held by the publishers listed in the PUBLISHER (PB) field (available

for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 20 Jul 2006 VOL 145 ISS 5
FILE LAST UPDATED: 20 Jul 2006 (20060720/ED)

New CAS Information Use Policies, enter HELP USAGETERMS for details.

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> d l12 cbib, ab 8,9,11,15,16,20,21

L12 ANSWER 8 OF 29 CA COPYRIGHT 2006 ACS on STN

132:52086 The physicochemical properties of some vegetable oils. Kuliev, R. Sh.; Shirinov, F. R.; Kuliev, F. A. (Volgograd Refinery, Russia). Chemistry and Technology of Fuels and Oils (Translation of Khimiya i Tekhnologiya Topliv i Masel), 35(4), 235-237 (English) 1999. CODEN: CTFOAK. ISSN: 0009-3092. Publisher: Consultants Bureau.

AB Physicochem. properties of several vegetable oils are presented and compared with those of crude petroleum for use as lubricating oils. Properties reported include d., viscosity, viscosity index, acid number, coking capacity, flash point, pour point, refractive index, color, saponification number, ash content, corrosivity.

L12 ANSWER 9 OF 29 CA COPYRIGHT 2006 ACS on STN

131:56440 Physicochemical characterization of oils from some Kenyan plants. Njagi, E. N. M.; Munyua, J. K.; Mark, A. G. (Department of Biochemistry, College of Health Sciences, University of Nairobi, Nairobi, Kenya). International Journal of BioChemPhysics, 6 & 7(1 & 2), 55-57 (English) 1998. CODEN: IJBOEY. ISSN: 1019-7648. Publisher: Centre for Nuclear Science Techniques, University of Nairobi.

AB The physiochem. characteristics of oils extracted from fruits and nuts collected from several parts of Kenya were determined. The parameters investigated included iodine value, acid value, saponification number, d., refractive index and viscosity. The iodine values were 78.05, 67.3, 83.13 and 6.98 for castor, green and purple avocado and coconut oils, resp. Acid values were between 0.840 and 2.244. Saponification nos. were 196.56, 261.80, 189.84 and 266.4 for castor, purple and green avocado and coconut oils, resp. D. values for all the oils were about the same and ranged between 0.8944 - 0.96027 gm/mL. Refractive index values were ND20, 1.4792, 1.4640, 1.4665 and ND60 1.4543 for castor, purple and green avocado oils and coconut oil, resp. Viscosities values ranged between 204.373 to 993.583 for castor and avocado oils while coconut oil was solid at room temperature. The data suggests that the oils from these Kenyan plants may be ideal for industrial use. Their cultivation should therefore be encouraged in order to minimize importation of similar oils.

L12 ANSWER 11 OF 29 CA COPYRIGHT 2006 ACS on STN

129:15515 Stable coconut cream alternative. Grant, Elizabeth R.; Norton, Ian Timothy; Foster, Timothy J.; Underdown, Jeffrey; Kimsey, Ian Michel (Unilever N.V., Neth.; Unilever PLC; Grant, Elizabeth R.; Norton, Ian Timothy; Foster, Timothy J.; Underdown, Jeffrey; Kimsey, Ian Michel). PCT Int. Appl. WO 9819553 A1 19980514, 28 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI,

SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG.

(English). CODEN: PIXXD2. APPLICATION: WO 1997-EP5921 19971021.

PRIORITY: EP 1996-307975 19961104.

AB Disclosed is sterilized, water-continuous fat containing emulsion comprising: 1-30 weight%, preferably 1-20 % of a vegetable or animal fat; 0.1-5 weight%, preferably 0.2-3 % of a protein; 0-2 weight% of an emulsifier composition; 0-10 weight% of a sweetener, in particular a carbohydrate; 0.01-2 weight% of a

flavor

composition; 0-1500 ppm. of cations; 0-5 weight% of a thickener composition; 0.01-5

weight% of stabilizer in particulate form; which composition has a viscosity at 30 °C of 1-200 mPa.s when measured at 50s-1 and after a storage period in the range of 1-36 wk at a temperature of 30 °C shows a creaming level in the range of 0-30 %. Such emulsions were found to be pourable and also to have a good storage stability, even for a period of up to 9 mo at high temps.

L12 ANSWER 15 OF 29 CA COPYRIGHT 2006 ACS on STN

124:200581 Mechanical properties of vegetable oils and fats. Alvarado, Juan de Dios (Facultad Ciencia Ingenieria Alimentos., Univ. Tecnica Ambato, Ecuador). Grasas y Aceites (Seville), 46(4-5), 264-9 (Spanish) 1995.

CODEN: GRACAN. ISSN: 0017-3495. Publisher: Instituto de la Grasa y sus Derivados.

AB Data of mech. properties are presented for crude oils from avocado pulp, lupine grain, peanuts, soybean, sesame, cotton, castor-oil, linseed, and passion fruit seeds; refined oils from sunflower, corn, peanut, olive and soybean; and cocoa, coconut, palm-oil, and kernel palm-oil fats.

Correlation equations which describe the effect of temperature on the refractive

index, d., viscosity, and surface tension are obtained, and

values of coefficient of expansion and activation energy for flow are calculated

L12 ANSWER 16 OF 29 CA COPYRIGHT 2006 ACS on STN

120:162092 Relationship between the molecular structure and deep fat frying properties of edible oils. Kimura, Masami; Harigaya, Hideko; Yanagisawa, Tamie; Takamura, Hitoshi; Matoba, Teruyoshi (Grad. Sch. Human Cult., Nara Women's Univ., Nara, 630, Japan). Nippon Kasei Gakkaishi, 44(12), 1027-32 (English) 1993. CODEN: NKGAE. ISSN: 0913-5227.

AB The relationship between the mol. structure and such deep fat frying properties of edible oils as oil absorption, dehydration, and oil drainage were investigated during deep fat frying by various monoacid-triacylglycerol mol. species and edible oils. As the carbon number of the fatty acid moieties of triacylglycerol increased, the initial rates of oil absorption and dehydration decreased; however, the maximal oil absorption and dehydration were similar. According to the oil drainage behavior, the oils were classified into two groups, i.e., solid fats and liquid oils. In the case of solid fats, the maximal oil drainage increased as the m.p. dropped. On the other hand, the oil drainage rate for liquid oils increased with decreasing viscosity. These results demonstrate that the mol. structure of edible oils affects the deep fat frying properties.

L12 ANSWER 20 OF 29 CA COPYRIGHT 2006 ACS on STN

111:59919 Interrelationships among the properties of fatty oils. Dutt, N. V. K.; Prasad, D. H. L. (Reg. Res. Lab., Hyderabad, 500 007, India). JAOCS, J. Am. Oil Chem. Soc., 66(5), 701-3 (English) 1989. CODEN: JJASDH.

AB Equations relating the properties of fatty oils, such as viscosity (η), iodine value (IV), and saponification value (SV), were developed, based on a model similar to the Antoine equation for vapor pressure. Two equations resulted. The first equation, $\log \eta = [-1.4 + 1.25 (IV/SV)] + [500 - 375(IV/SV)] / [(t + 140) - 85(IV/SV)]$ (t = temperature in °C), gave an average absolute deviation of 13.0% at 77 data points of several fatty oils. The second equation, $\log \eta = 0.6298 + [273.660(t + 88.81)]$,

gave an average absolute deviation of 14.5% at 89 data points of a larger group of fatty oils ranging from almond to tallow. The two equations could be used conveniently to predict either viscosity, iodine value, or saponification value, when the other two properties were known, for design purposes.

L12 ANSWER 21 OF 29 CA COPYRIGHT 2006 ACS on STN

105:23288 Rheological properties of emulsified fats with regard to their optimum consistency. Stern, Petr; Pokorny, Jan; Dobiasova, Stanislava; Davidek, Jiri; Cmolik, Jiri (Ustav Hydrodyn., CSAV, Prague, Czech.). Prumysl Potravin, 37(1), 19-21 (Czech) 1986. CODEN: PPOTAP. ISSN: 0033-1988.

AB Basic rheol. characteristics (static and dynamic yield point and apparent viscosity) and sensory characteristics (hardness determined by cutting and spreadability with a knife) were determined at 5-25° for 2 types of com. edible emulsified fats: (1) a common semi-soft fat containing vegetable oil 30, coconut oil 10, and hydrogenated vegetable oil 60% and (2) a soft fat containing 60% vegetable oil and 40% hydrogenated vegetable oil. High correlation coeffs. were obtained between the rheol. and sensory characteristics indicating optimum properties of both types of fats. The optimum consistency was observed in the case of soft fat at lower temps. than in the case of semi-soft fat.

=> d 113 cbib,ab 17,23,28,31,32,33,39,47,51,53,60,67,100,113,128,130,134

L13 ANSWER 17 OF 134 CA COPYRIGHT 2006 ACS on STN

139:35478 Comparative studies on physical properties of vegetable oils and their blends after frying. Susheelamma, N. S.; Asha, M. R.; Ravi, R.; Kumar, A. K. Vasanth (Department of Sensory Science, Central Food Technological Research Institute, Mysore, 570013, India). Journal of Food Lipids, 9(4), 259-276 (English) 2002. CODEN: JFFLES. ISSN: 1065-7258. Publisher: Food & Nutrition Press, Inc..

AB Phys. properties of six commonly used oils and three blends consisting of three oils in each blend were studied after three successive frying of 'poories' (fried snack from fattened dough of refined wheat flour). The changes in viscosity, CIE trans-reflectance color and related parameters, UV-Visible spectra and UV-spectra of oil samples in solvent system (chloroform:methanol; 2:1, volume/volume) were studied. The results showed that viscosity and color of the oils changed to a much higher extent after first frying than subsequent fryings. The hue angle followed a similar trend. Changes in the UV-spectra in the solvent system indicated an increase in the formation of conjugated compds. after successive fryings. Peroxide values (PV) also increased after frying. Principal Component Anal. (PCA) plots of the data indicated that among oils examined groundnut oil and soy oil in combination with other oils were preferred for frying. Use of small amts. of unrefined oils (filtered) such as mustard oil or sesame oil which have a high content of natural antioxidants was beneficial as formation of conjugated compds. and increase in peroxide value was minimized after successive frying using blended oils.

L13 ANSWER 23 OF 134 CA COPYRIGHT 2006 ACS on STN

135:106667 Thickened oil compositions of edible oil. Eini, Meir; Tamarkin, Dov (Thixo Ltd., Israel). PCT Int. Appl. WO 2001050873 A1 20010719, 118 pp. DESIGNATED STATES: W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, CY, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG, TR. (English). CODEN: PIXXD2. APPLICATION: WO 2001-IL24 20010110. PRIORITY: IL 2000-133968 20000110; IL 2000-133969

20000110; US 2000-526509 20000316; IL 2000-137051 20000627; IL 2000-137052 20000627; US 2000-216162P 20000703; US 2000-653267 20000831.

- AB Thickened, preferably thixotropic oil compns. are disclosed comprised of high proportions of edible oils, in particular unsatd. oils, and edible solidifying agents, in particular long-chain fatty acids and/or long-chain fatty alcs., for use in the preparation of edible foods and medicinal/therapeutic products. Further disclosed are methods for the production of the thickened, thixotropic oil compns., edible food, and medicinal/therapeutic products containing same and methods of their production and consumption.

L13 ANSWER 28 OF 134 CA COPYRIGHT 2006 ACS on STN

132:193459 Density and viscosity of vegetable oils. Rodenbush, C. M.; Hsieh, F. H.; Viswanath, D. S. (Departments of Chemical Engineering, University of Missouri-Columbia, Columbia, MO, 65211, USA). Journal of the American Oil Chemists' Society, 76(12), 1415-1419 (English) 1999. CODEN: JAOCA7. ISSN: 0003-021X. Publisher: AOCS Press.

- AB A generalized method was developed to estimate the liquid d. of vegetable oils and fatty acids. The correlation for vegetable oils was based on fatty acid critical properties and composition of the oil. The correlations predicted the d. of vegetable oils and fatty acids with an average absolute deviation of 0.21 and 0.77%, resp. This method is slightly more accurate in predicting vegetable oil d. and simpler than the method of J. D. Halvorsen et al. (1993). A method is introduced that predicts viscosity from d. data, thus relating two key properties of vegetable oils.

L13 ANSWER 31 OF 134 CA COPYRIGHT 2006 ACS on STN

132:52086 The physicochemical properties of some vegetable oils. Kuliev, R. Sh.; Shirinov, F. R.; Kuliev, F. A. (Volgograd Refinery, Russia). Chemistry and Technology of Fuels and Oils (Translation of Khimiya i Tekhnologiya Topliva i Masel), 35(4), 235-237 (English) 1999. CODEN: CTFOAK. ISSN: 0009-3092. Publisher: Consultants Bureau.

- AB Physicochem. properties of several vegetable oils are presented and compared with those of crude petroleum for use as lubricating oils. Properties reported include d., viscosity, viscosity index, acid number, coking capacity, flash point, pour point, refractive index, color, saponification number, ash content, corrosivity.

L13 ANSWER 32 OF 134 CA COPYRIGHT 2006 ACS on STN

131:89296 Enzymic interesterification of blends of castor oil and some oils rich in saturated fatty acids. Ghosh, Mahua; Bhattacharyya, Dipak K. (Department Chemical Technology, Oil Technology Division, Calcutta Univ., Calcutta, 700009, India). Fett/Lipid, 101(6), 214-216 (English) 1999. CODEN: FELIFX. ISSN: 0931-5985. Publisher: Wiley-VCH Verlag GmbH.

- AB Interesterification of castor oil blended with some oils rich in saturated fatty acids was done with the help of 1,3-specific lipase from *Mucor miehei* in order to alter its viscosity characteristics and adhesion properties by the introduction of saturated fatty acid mols. The interesterification was done by an oil blend ratio of 50:50. 10% Enzyme were used. Temperature was kept at 60° under 2-5 mm Hg pressure with constant stirring, and the reactions were carried out for 6 h. The products were filtered to remove the enzyme and then analyzed for slip point, sp. gr., and kinematic viscosity. The slip point of the interesterified products was found to be much lower than the parent blend and was in the range of 15-25°. Sp. gr. and iodine value of the products were in comparison with the theor. ones. A very large depression in kinematic viscosity was found with every interesterified product from original castor oil and also from the blends at 3 different temps.

L13 ANSWER 33 OF 134 CA COPYRIGHT 2006 ACS on STN

131:56440 Physicochemical characterization of oils from some Kenyan plants.

Njagi, E. N. M.; Munyua, J. K.; Mark, A. G. (Department of Biochemistry, College of Health Sciences, University of Nairobi, Nairobi, Kenya). International Journal of BioChemiPhysics, 6 & 7(1 & 2), 55-57 (English) 1998. CODEN: IJBOEY. ISSN: 1019-7648. Publisher: Centre for Nuclear Science Techniques, University of Nairobi.

- AB The physiochem. characteristics of oils extracted from fruits and nuts collected from several parts of Kenya were determined. The parameters investigated included iodine value, acid value, saponification number, d., refractive index and viscosity. The iodine values were 78.05, 67.3, 83.13 and 6.98 for castor, green and purple avocado and coconut oils, resp. Acid values were between 0.840 and 2.244. Saponification nos. were 196.56, 261.80, 189.84 and 266.4 for castor, purple and green avocado and coconut oils, resp. D. values for all the oils were about the same and ranged between 0.8944 - 0.96027 gm/mL. Refractive index values were ND20, 1.4792, 1.4640, 1.4665 and ND60 1.4543 for castor, purple and green avocado oils and coconut oil, resp. Viscosities values ranged between 204.373 to 993.583 for castor and avocado oils while coconut oil was solid at room temperature. The data suggests that the oils from these Kenyan plants may be ideal for industrial use. Their cultivation should therefore be encouraged in order to minimize importation of similar oils.

L13 ANSWER 39 OF 134 CA COPYRIGHT 2006 ACS on STN

129:15515 Stable coconut cream alternative. Grant, Elizabeth R.; Norton, Ian Timothy; Foster, Timothy J.; Underdown, Jeffrey; Kimsey, Ian Michel (Unilever N.V., Neth.; Unilever PLC; Grant, Elizabeth R.; Norton, Ian Timothy; Foster, Timothy J.; Underdown, Jeffrey; Kimsey, Ian Michel). PCT Int. Appl. WO 9819553 A1 19980514, 28 pp. DESIGNATED STATES: W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM; RW: AT, BE, BF, BJ, CF, CG, CH, CI, CM, DE, DK, ES, FI, FR, GA, GB, GR, IE, IT, LU, MC, ML, MR, NE, NL, PT, SE, SN, TD, TG. (English). CODEN: PIXXD2. APPLICATION: WO 1997-EP5921 19971021. PRIORITY: EP 1996-307975 19961104.

- AB Disclosed is sterilized, water-continuous fat containing emulsion comprising: 1-30 weight%, preferably 1-20 % of a vegetable or animal fat; 0.1-5 weight%, preferably 0.2-3 % of a protein; 0-2 weight% of an emulsifier composition; 0-10 weight% of a sweetener, in particular a carbohydrate; 0.01-2 weight% of a flavor composition; 0-1500 ppm. of cations; 0-5 weight% of a thickener composition; 0.01-5 weight% of stabilizer in particulate form; which composition has a viscosity at 30 °C of 1-200 mPa.s when measured at 50s-1 and after a storage period in the range of 1-36 wk at a temperature of 30 °C shows a creaming level in the range of 0-30 %. Such emulsions were found to be pourable and also to have a good storage stability, even for a period of up to 9 mo at high temps.

L13 ANSWER 47 OF 134 CA COPYRIGHT 2006 ACS on STN

124:200581 Mechanical properties of vegetable oils and fats. Alvarado, Juan de Dios (Facultad Ciencia Ingenieria Alimentos., Univ. Tecnica Ambato, Ecuador). Grasas y Aceites (Seville), 46(4-5), 264-9 (Spanish) 1995. CODEN: GRACAN. ISSN: 0017-3495. Publisher: Instituto de la Grasa y sus Derivados.

- AB Data of mech. properties are presented for crude oils from avocado pulp, lupine grain, peanuts, soybean, sesame, cotton, castor-oil, linseed, and passion fruit seeds; refined oils from sunflower, corn, peanut, olive and soybean; and cocoa, coconut, palm-oil, and kernel palm-oil fats. Correlation equations which describe the effect of temperature on the refractive index, d., viscosity, and surface tension are obtained, and values of coefficient of expansion and activation energy for flow are calculated.

L13 ANSWER 51 OF 134 CA COPYRIGHT 2006 ACS on STN

120:162092 Relationship between the molecular structure and deep fat frying properties of edible oils. Kimura, Masami; Harigaya, Hideko; Yanagisawa, Tamie; Takamura, Hitoshi; Matoba, Teruyoshi (Grad. Sch. Human Cult., Nara Women's Univ., Nara, 630, Japan). Nippon Kasei Gakkaishi, 44(12), 1027-32 (English) 1993. CODEN: NKGAE. ISSN: 0913-5227.

AB The relationship between the mol. structure and such deep fat frying properties of edible oils as oil absorption, dehydration, and oil drainage were investigated during deep fat frying by various monoacid-triacylglycerol mol. species and edible oils. As the carbon number of the fatty acid moieties of triacylglycerol increased, the initial rates of oil absorption and dehydration decreased; however, the maximal oil absorption and dehydration were similar. According to the oil drainage behavior, the oils were classified into two groups, i.e., solid fats and liquid oils. In the case of solid fats, the maximal oil drainage increased as the m.p. dropped. On the other hand, the oil drainage rate for liquid oils increased with decreasing viscosity. These results demonstrate that the mol. structure of edible oils affects the deep fat frying properties.

L13 ANSWER 53 OF 134 CA COPYRIGHT 2006 ACS on STN

118:41153 Viscosities of vegetable oils and fatty acids. Nouredдини, H.; Teoh, B. C.; Clements, L. Davis (Dep. Chem. Eng., Univ. Nebraska, Lincoln, NE, 68588-0126, USA). Journal of the American Oil Chemists' Society, 69(12), 1189-91 (English) 1992. CODEN: JAOCA7. ISSN: 0003-021X.

AB Data for viscosity as a function of temperature from 24 to 110° were measured for a number of vegetable oils (crambe, rapeseed, corn, soybean, milkweed, coconut, and lesquerella) and 8 fatty acids in the range from C9 to C22. The viscosity measurements were performed according to ASTM test methods D 445 and D 446. Several correlations were fitted to the exptl. data. Correlation consts. for the best fit are presented. The correlation consts. are valuable for designing or evaluating such chemical process equipment as heat exchangers, reactors, distillation columns, mixing vessels, and process piping.

L13 ANSWER 60 OF 134 CA COPYRIGHT 2006 ACS on STN

111:59919 Interrelationships among the properties of fatty oils. Dutt, N. V. K.; Prasad, D. H. L. (Reg. Res. Lab., Hyderabad, 500 007, India). JAOCS, J. Am. Oil Chem. Soc., 66(5), 701-3 (English) 1989. CODEN: JJASDH.

AB Equations relating the properties of fatty oils, such as viscosity (η), iodine value (IV), and saponification value (SV), were developed, based on a model similar to the Antoine equation for vapor pressure. Two equations resulted. The first equation, $\log \eta = [-1.4 + 1.25 (IV/SV)] + [500 - 375 (IV/SV)] / [(t + 140) - 85 (IV/SV)]$ (t = temperature in °C), gave an average absolute deviation of 13.0% at 77 data points of several fatty oils. The second equation, $\log \eta = 0.6298 + [273.660(t + 88.81)]$, gave an average absolute deviation of 14.5% at 89 data points of a larger group of fatty oils ranging from almond to tallow. The two equations could be used conveniently to predict either viscosity, iodine value, or saponification value, when the other two properties were known, for design purposes.

L13 ANSWER 67 OF 134 CA COPYRIGHT 2006 ACS on STN

102:130494 Physical properties of oils and mixtures of oils. Timms, R. E. (Kempas Edible Oil Sendirian Berhad, Johore, Malay.). JAOCS, J. Am. Oil Chem. Soc., 62(2), 241-8 (English) 1985. CODEN: JJASDH.

AB A review with 40 refs. on the d., sp. heat, heat of fusion, and viscosity of palm, palm kernel, and coconut oils and their mixts.

L13 ANSWER 100 OF 134 CA COPYRIGHT 2006 ACS on STN

71:103225 Continuous viscosity measurement in stirred vessels.

Kleinschmidt, Ernst (Herbol-Werke, Herbig-Haarhaus A.-G., Cologne, Fed.

Rep. Ger.). Paint Technology, 33(9), 34-6, 39-44 (English) 1969. CODEN: PATEA2. ISSN: 0030-9524.

AB A method is described which enables the viscosity of resin melts in stirred vessels to be measured continuously during manufacture. The method used a blade of magnetostrictive material vibrating at an ultrasonic frequency; damping of the blade was dependent on the viscosity of the liquid. Calibration curves of melt viscosity and solution viscosity were prepared by taking resin samples of different solution viscosities from the reactor and determining their melt viscosities. Resins used were a nondrying alkyd, a linseed oil alkyd, and an epoxide ester of dehydrated ricinoleic acid. In continuous measurement the turbulent flow, which causes difficulties in measurements, was converted to a laminar flow. The lower part of the probe was bent at right angles and surrounded by a guide tube which tapered towards the end of the probe. Viscosity measurements were carried out by means of the new probe on coconut oil, linseed oil, and soybean oil alkyd resins.

L13 ANSWER 113 OF 134 CA COPYRIGHT 2006 ACS on STN
63:12947 Original Reference No. 63:2308e-f The stability of oils and fats after prolonged storage in retail packs. Winter, E. (Bundesforschungsanstalt Lebensmittelfrischhaltung, Karlsruhe, Germany). Fette, Seifen, Anstrichmittel, 67(2), 124-30 (German) 1965. CODEN: FSASAX. ISSN: 0015-038X.

AB The storage properties of peanut oil, soybean oil, sunflower seed oil, and coconut oil in packages used by the trade, were studied at 0-25°. As criteria of stability, the peroxide, thiobarbituric acid, and acid values were compared with the taste, odor, and viscosity of the samples over a period of 3 years. The oxidative deterioration can be considerably reduced by sealing the containers under N or in vacuo.

L13 ANSWER 128 OF 134 CA COPYRIGHT 2006 ACS on STN
34:44676 Original Reference No. 34:6838a-b The physical properties of mixtures of coconut and castor oil. Clemente, Amando; Rillo, Socorro Univ. Philippines Nat. and Applied Sci. Bull., 7, 1319-25 (Unavailable) 1940.

AB Solns. of castor oil in coconut oil were compared with Castrol "XL" and Mobiloil "A" (cf. Medina and C., C. A. 28, 7047.6). The concentration of castor oil ranged from 5% to 100%, in steps of 5%. The sp. gr.,

surface tension and viscosity were measured at temps. between 30° and 95° in 5° intervals. The solidifying point was also measured. n was determined at temps. from 25° to 50° in 5° intervals. Phys. properties indicate a mixture of castor oil and coconut oil may possibly be used as a substitute for Castrol "XL" or Mobiloil "A" as a lubricating oil. 17 references.

L13 ANSWER 130 OF 134 CA COPYRIGHT 2006 ACS on STN
28:57808 Original Reference No. 28:7047f-h Physical properties of some Philippine vegetable oils. Medina, Florencio A.; Clemente, Amando (Univ. Philippines Natural and Applied Sci.). Bull., 4(No. 1), 61-91 (Unavailable) 1934.

AB The oils studied are: calumpang, cashew, coconut, kapok, lumbang, palo-maria de la playa, peanut, pili-nut, tangan-tangan, tuba, Castrol XL and Mobiloil A. Their colors, optical rotations and melting and congealing points are tabulated. Only palo-maria and tangan-tangan oils show appreciable rotation. A low congealing point seems to indicate a drying oil. All the oils are completely soluble in Et₂O, C₆H₆, benzine, Me₂CO, CHCl₃, CCl₄ and CS₂. Only tangan-tangan oil is completely soluble in 95% EtOH even at room temperature; this is probably due to a high percentage of hydroxylated fatty acids and can be used in identifying this oil. Tables and graphs show the variation of sp. gr., n, surface tension and viscosity with temperature at 5° intervals. If Castrol XL and

Mobiloil A are taken as standards, calumpang oil has the viscosity required of a lubricating oil.

L13 ANSWER 134 OF 134 CA COPYRIGHT 2006 ACS on STN

14:12829 Original Reference No. 14:2387a-c Blended fatty food products.
Schwarcman, A. US 1342827 19200608 (Unavailable). APPLICATION: US .

AB Blended fatty food products are prepared of a predetd. hardness or viscosity by forming an emulsion containing separated particles of various oils or fats of selected different m. p. A product suitable for cooking purposes may be formed as follows: Peanut oil is warmed to about 45° and thoroughly mixed with 5-8% H2O so as to reduce the oil to very small particles. The H2O may carry a small amount of an emulsifying agent such as gelatin, Irish moss, Iceland moss, casein or similar substances. After emulsification of the oil, it is mixed with 15-30% of beef stearin or other similar fat of relatively high m. p. (The percentages given are based on the total weight of the final product.) Cottonseed oil, palm oil, lard oil, butter oil or refined coconut oil may be used in making up similar products. A product resembling butter in most of its properties may be thus formed by churning together suitably selected ingredients.

=> log y

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
63.28	319.27

FULL ESTIMATED COST

DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)

SINCE FILE	TOTAL
ENTRY	SESSION
-17.04	-23.43

CA SUBSCRIBER PRICE

STN INTERNATIONAL LOGOFF AT 09:28:15 ON 28 JUL 2006